



Periodic Solutions of Coupled Nonlinear Schrödinger Equations in Nonlinear Optics: the Nonresonant Case

B. GRÉBERT

M.I.P. U.M.R. 9974 C.N.R.S., Université Paul Sabatier, Toulouse III
118, route de Narbonne, 31062 Toulouse Cedex, France

J. C. GUILLOT

Département de Mathématiques, U.R.A. C.N.R.S. 742
Institut Galilée - Université Paris-Nord, 93430 Villetaneuse, France
and

C.M.L.A., E.N.S. Cachan, U.R.A. C.N.R.S. 1661
61, avenue du Président Wilson, 94235 Cachan Cedex, France

(Received March 1995; accepted May 1995)

Abstract—We construct periodic solutions to coupled nonlinear one-dimensional Schrödinger equations with periodic boundary conditions in the nonresonant case.

Keywords—Periodic Schrödinger equations, Nash-Moser method.

The purpose of this note is to construct periodic solutions of coupled nonlinear one-dimensional Schrödinger equations with periodic boundary conditions. In particular, these equations are those for the dimensionless envelopes of the electromagnetic fields (u, v) in nonlinear birefringent fibers [1,2]. Precisely, we consider the following equations:

$$\begin{aligned} i \frac{\partial u}{\partial t} &= -\frac{\partial^2 u}{\partial x^2} + q_1(x) u + \alpha(x) v + g_1(u, \bar{u}, v, \bar{v}, x), \\ i \frac{\partial v}{\partial t} &= -\frac{\partial^2 v}{\partial x^2} + q_2(x) v + \alpha(x) u + g_2(u, \bar{u}, v, \bar{v}, x) \end{aligned} \quad (1)$$

with $u(x + \pi, t) = u(x)$ and $v(x + \pi, t) = v(x, t)$.

We proceed along the lines of the method recently developed by Craig and Wayne to construct time periodic solutions to the nonlinear wave and Schrödinger equations [3,4]. The results of Craig and Wayne have been recently generalized by Bourgain who has been able to construct quasi-periodic solutions of 1-D nonlinear wave and Schrödinger equations and 2-D nonlinear Schrödinger equations [5,6]. Craig and Wayne follow the Lyapunov-Schmidt argument which consists in splitting the problem in a resonant finite dimensional piece given by the Q -equation and the remainder of the problem, the P -equation, which is infinite dimensional and contains small divisors issues. We suppose that

$$\begin{aligned} g_1(u, \bar{u}, v, \bar{v}, x) &= \partial_{\bar{u}} G(u, \bar{u}, v, \bar{v}, x), \\ g_2(u, \bar{u}, v, \bar{v}, x) &= \partial_{\bar{v}} G(u, \bar{u}, v, \bar{v}, x), \end{aligned} \quad (2)$$

where $G(u, \bar{u}, v, \bar{v}, x)$ is a real-valued function.

In nonlinear optics, a typical example is

$$G(u, \bar{u}, v, \bar{v}, x) = \frac{1}{2}|u|^4 + a(x)|u|^2|v|^2 + \frac{1}{2}.$$

The function $G(z_1, z_2, z_3, z_4, x)$ is supposed to be periodic in x with period π , analytic in the domain $\{(z_1, z_2, z_3, z_4, x) \in \mathbb{C}^5 \mid |z_i| < 1, |\operatorname{Im} x| < \sigma\}$, and

$$G(z_1, z_2, z_3, z_4, x) = \sum_{|\alpha| \geq 4} a_\alpha(x) z_1^{\alpha_1} z_2^{\alpha_2} z_3^{\alpha_3} z_4^{\alpha_4}, \quad (3)$$

with $\alpha = (\alpha_1, \alpha_2, \alpha_3, \alpha_4) \in \mathbb{N}^4$ and $|\alpha| = \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4$. Thus, g_1 and g_2 are cubic in u, \bar{u}, v, \bar{v} .

The set of all functions G satisfying these hypotheses with the topology of uniform convergence is denoted by \mathcal{A}_σ . We also assume that the potentials $q_1(x)$, $q_2(x)$ and $a(x)$ are periodic in x with period π , analytic in the domain $\{x \in \mathbb{C} \mid |\operatorname{Im} x| < \sigma\}$. The set of all potentials satisfying these hypotheses again with the topology of uniform convergence is denoted by \mathcal{Q}_σ .

CASE A. Let us first consider the case where $\alpha \equiv 0$ and $q_1 \neq q_2$. Let $(\psi_j(x))_{j \geq 0}$ (respectively, $(\tilde{\psi}_j(x))_{j \geq 0}$) be the normalized eigenfunctions of $-\frac{d^2}{dx^2} + q_1(x)$ (respectively, $-\frac{d^2}{dx^2} + q_2(x)$) with periodic boundary conditions associated with the corresponding eigenvalues $(w_j)_{j \geq 0}$ (respectively, $(\tilde{w}_j)_{j \geq 0}$).

Rescaling the time variable in (1), i.e., setting $\xi = t\Omega$ where Ω is a frequency, we now consider

$$\begin{aligned} -i\Omega \frac{\partial u}{\partial \xi} &= -\frac{\partial^2 u}{\partial x^2} + q_1(x)u + g_1(u, \bar{u}, v, \bar{v}, x), \\ -i\Omega \frac{\partial v}{\partial \xi} &= -\frac{\partial^2 v}{\partial x^2} + q_1(x)v + g_2(u, \bar{u}, v, \bar{v}, x). \end{aligned} \quad (4)$$

Set

$$\Phi_{2j}(x) = \begin{pmatrix} \psi_j(x) \\ 0 \end{pmatrix}, \quad \Phi_{2j+1}(x) = \begin{pmatrix} 0 \\ \tilde{\psi}_j(x) \end{pmatrix}.$$

Looking for a periodic solution $\begin{pmatrix} u(x, \xi) \\ v(x, \xi) \end{pmatrix}$ to (4) we expand it in an eigenfunction expansion

$$\begin{pmatrix} u(x, \xi) \\ v(x, \xi) \end{pmatrix} = \sum_{(j,k) \in \mathbb{N} \times \mathbb{Z}} \hat{u}(j, k) \Phi_j(x) e^{-ik\xi}. \quad (5)$$

Insert (5) into (4) to get the nonlinear equation

$$W(\hat{u})(j, k) + V(\Omega) \hat{u}(j, k) = 0, \quad j \in \mathbb{N}, \quad k \in \mathbb{Z} \quad (6)$$

with

$$\begin{aligned} V(\Omega) \hat{u}(2j, k) &= (w_j - k\Omega) \hat{u}(2j, k), \\ V(\Omega) \hat{u}(2j+1, k) &= (\tilde{w}_j - k\Omega) \hat{u}(2j+1, k) \end{aligned}$$

and

$$W(\hat{u})(j, k) = \int_0^\pi \int_0^{2\pi} \left\langle \begin{pmatrix} g_1(u, \bar{u}, v, \bar{v}, x) \\ g_2(u, \bar{u}, v, \bar{v}, x) \end{pmatrix}, \Phi_j(x) e^{-ik\xi} \right\rangle_{\mathbb{C}^2} dx d\xi,$$

with $W(T_\theta \hat{u}) = T_\theta W(\hat{u})$, where $(T_\theta \hat{v})(j, k) = e^{ik\theta} \hat{v}(j, k)$.

Equation (6) is split into a finite dimensional and an infinite dimensional part according to the Lyapunov-Schmidt procedure.

Fix $j_0 \in \mathbb{N}$ and set

$$\mathcal{N} = \{(j, k) \in \mathbb{N} \times \mathbb{Z} \mid V(w_{j_0})(j, k) = 0\}. \quad (7)$$

Generically, for $q_1, q_2 \in \mathcal{Q}_\sigma$, and $G \in \mathcal{A}_\sigma$, $\dim N = 1$ [3] and from now on, we suppose that $N = \{(2j_0, 1)\}$. We could have fixed \tilde{w}_{j_0} as well.

Given N , let Q be the orthogonal projection onto $\ell^2(N) \subset \ell^2(\mathbb{N} \times \mathbb{Z})$ and set $P = Q^\perp$. Equation (6) is equivalent to

$$PW(\hat{u})(j, k) + PV(\Omega)\hat{u}(j, k) = 0, \quad (8)$$

$$QW(\hat{u})(j, k) + QV(\Omega)\hat{u}(j, k) = 0. \quad (9)$$

Equation (8) is solved by using a Nash-Moser implicit function theorem and (9) by using bifurcation theory. In order to apply a Nash-Moser procedure, it is necessary to introduce a family of Hilbert subspaces of $\ell^2(\mathbb{N} \times \mathbb{Z})$, i.e., the spaces denoted by $H_{m,s}$, $m \geq 0$, $s \geq 0$ and defined in terms of the $\hat{u}(j, k)$'s as in [4]. Furthermore, we need to satisfy nonresonance conditions. Thus, we suppose that the two sequences $(w_j)_{\substack{j \geq 0 \\ j \neq j_0}}$ and $(\tilde{w}_j)_{j \geq 0}$ are (d_0, L_0) nonresonant with \tilde{w}_{j_0} , i.e., that, for every $(j, k) \in B(L_0) \setminus N$, we have

$$|kw_{j_0} - w_j| > d_0, \quad |kw_{j_0} - \tilde{w}_j| > d_0, \quad (10)$$

with L_0 sufficiently large and d_0 depending in L_0 in an appropriate way (cf. [4]). Here, $B(L_0) = \{(j, k) \mid j + |k| \leq L_0\}$ and we suppose that $N \subset B(L_0)$.

Solutions to (8) are constructed by perturbing the solution $\varphi(p) = p\delta_{\{(j,k),(j_0,1)\}}$ of the linear part with $p \in \mathbb{C}$. A major part is devoted to the analysis of the linearized operator $PV(\Omega) + DPW(\hat{u})$. Inversion of linear operators proceeds along the lines of the method developed by Fröhlich and Spencer [7]. The only condition that the nonlinear part of (9) must satisfy is the so-called twist condition

$$\left. \frac{\partial^3}{\partial r^3} W(\varphi(r)) \right|_{r=0} (2j_0, 1) \neq 0. \quad (11)$$

This condition is used to apply the Morse Lemma.

According to (3), the twist condition is

$$\int_0^\pi a_{2,2,0,0}(x) |\psi_{j_0}(x)|^4 dx \neq 0.$$

Fixing \tilde{w}_{j_0} , we obtain

$$\int_0^\pi a_{0,0,2,2}(x) |\tilde{\psi}_{j_0}(x)|^4 dx \neq 0.$$

We finally get the following theorem.

THEOREM 1. *Consider (1) with $\alpha = 0$, $q_1 \neq q_2$ and periodic boundary conditions. Consider w_{j_0} . There exists an open dense set $\mathcal{D}_{j_0} \subset \mathcal{A}_\sigma \times \mathcal{Q}_\sigma \times \mathcal{Q}_\sigma$ such that, if $(G, q_1, q_2) \in \mathcal{D}_{j_0}$, equation (1) has time periodic solutions close to $\Phi_{2j_0}(x) e^{i w_{j_0} t}$.*

More precisely, there exists $r_0 > 0$, a C^3 function $\Omega(r)$ defined on $[0, r_0)$ and a Cantor set $C \subset [0, r_0)$ of positive measure, such that, if $r \in C$, there exist time periodic solutions $u(x, t; r)$ and $v(x, t; r)$ of (1) with frequency $\Omega(r)$ such that

$$\left| u(x, t; r) - r \psi_{j_0}(x) e^{i \Omega(r) t} \right| \leq C r^2, \quad |v(x, t; r)| \leq C r^2,$$

and

$$|\Omega(r) - w_{j_0}| \leq C r^2.$$

REMARKS.

- (i) The same statement is true if we consider \tilde{w}_{j_0} instead of w_{j_0} with obvious changes.
- (ii) $\{\Omega(r) \mid r \in C\}$ is also a Cantor set and $0 \in C$.
- (iii) Twist conditions are partly used to show that the measure of C (respectively, \tilde{C}) is greater than $r_0/2$.

CASE B. We now consider (1) in the case where $\alpha \neq 0$ and $q_1 = q_2 = q$ with $\alpha, q \in \mathcal{Q}_\sigma$. We then have the following system:

$$\begin{aligned} i \frac{\partial u}{\partial t} + \frac{\partial^2 u}{\partial x^2} &= q(x) u + \alpha(x) v + g_1(u, \bar{u}, v, \bar{v}, x), \\ i \frac{\partial v}{\partial t} + \frac{\partial^2 v}{\partial x^2} &= q(x) v + \alpha(x) u + g_2(u, \bar{u}, v, \bar{v}, x). \end{aligned} \quad (12)$$

Set

$$U(x, t) = u(x, t) + v(x, t) \quad \text{and} \quad V(x, t) = u(x, t) - v(x, t).$$

Then, (12) is equivalent to

$$\begin{aligned} i \frac{\partial U}{\partial t} + \frac{\partial^2 U}{\partial x^2} &= (q(x) + \alpha(x)) U + \tilde{g}_1(U, \bar{U}, V, \bar{V}, x), \\ i \frac{\partial V}{\partial t} + \frac{\partial^2 V}{\partial x^2} &= (q(x) - \alpha(x)) V + \tilde{g}_2(U, \bar{U}, V, \bar{V}, x), \end{aligned} \quad (13)$$

with $\tilde{g}_1 = \partial_{\bar{U}} \tilde{G}$, $\tilde{g}_2 = \partial_{\bar{V}} \tilde{G}$ and

$$\tilde{G}(U, \bar{U}, V, \bar{V}, x) = G(u, \bar{u}, v, \bar{v}, x).$$

Therefore, we can now apply Theorem 1.

Let $(\varphi_j(x))_{j \geq 0}$ (respectively, $(\tilde{\varphi}_j(x))_{j \geq 0}$) be the normalized eigenfunctions of $-\frac{d^2}{dx^2} + q + \alpha$ (respectively, $-\frac{d^2}{dx^2} + q - \alpha$) with periodic boundary conditions associated with the corresponding eigenvalues $(\lambda_j)_{j \geq 0}$ (respectively, $(\tilde{\lambda}_j)_{j \geq 0}$).

Consider λ_{j_0} . Generically, for $(G, q, \alpha) \in \mathcal{A}_\sigma \times \mathcal{Q}_\sigma \times \mathcal{Q}_\sigma$, $\dim N = \{(j_0, 1)\}$. Suppose that $(\lambda_j)_{\substack{j \neq j_0 \\ j \geq 0}}$ and $(\tilde{\lambda}_j)_{j \geq 0}$ satisfy nonresonance conditions with respect to λ_{j_0} as in part A. We then have the following theorem.

THEOREM 2. *Consider (12) with periodic boundary conditions. Consider λ_{j_0} . There exists an open dense set $\mathcal{F}_{j_0} \subset \mathcal{A}_\sigma \times \mathcal{Q}_\sigma \times \mathcal{Q}_\sigma$ such that, if $(G, \alpha, q) \in \mathcal{F}_{j_0}$, then (12) has time periodic solutions close to $\begin{pmatrix} \varphi_{j_0}(x) e^{i\lambda_{j_0}t} \\ \varphi_{j_0}(x) e^{i\lambda_{j_0}t} \end{pmatrix}$. Thus, there exists $\rho_0 > 0$, a C^3 function $F(\rho)$ defined on $[0, \rho_0)$ and a Cantor set $K \subset [0, \rho_0)$ of positive measure, such that, if $\rho \in K$, there exist time periodic solutions $u(x, t; \rho)$ and $v(x, t; \rho)$ of (12) with frequency $F(\rho)$ such that*

$$\begin{aligned} |u(x, t; \rho) - \rho \varphi_{j_0}(x) e^{iF(\rho)t}| &\leq C \rho^2, \\ |v(x, t; \rho) - \rho \varphi_{j_0}(x) e^{iF(\rho)t}| &\leq C \rho^2, \\ |F(\rho) - \lambda_{j_0}| &\leq C \rho^2. \end{aligned}$$

Remarks following Theorem 1 are still true for Theorem 2. If we consider $\tilde{\lambda}_{j_0}$ instead of λ_{j_0} , time periodic solutions to (12) are close to $\begin{pmatrix} \varphi_{j_0}(x) \\ -\varphi_{j_0}(x) \end{pmatrix} e^{i\tilde{\lambda}_{j_0}t}$.

REFERENCES

1. A.C. Newell and J.V. Moloney, *Nonlinear Optics*, Addison-Wesley, (1992).
2. F. Abdullaev, S. Darmanyan and P. Khabibullaev, Optical solitons, In *Springer Series in Nonlinear Dynamics*, (1993).
3. W. Craig and C. Wayne, Newton's method and periodic solutions of nonlinear wave equations, *C.P.A.M.* **46**, 1409–1501 (1993).
4. W. Craig and C. Wayne, Periodic solutions of nonlinear Schrödinger equations and the Nash-Moser method, preprint.
5. J. Bourgain, Construction of quasi-periodic solutions of Hamiltonian perturbations of linear equations and applications to nonlinear P.D.E., Preprint, *I.H.E.S.* (August 1994); *I.M.R.N.* **11** (to appear).
6. J. Bourgain, Quasi-periodic solutions of Hamiltonians perturbations of 2D linear Schrödinger equations, Preprint *I.H.E.S.*, (January 1995).
7. J. Fröhlich and T. Spencer, Absence of diffusion in the Anderson tight binding model for large disorder or low energy, *Comm. Math. Phys.* **88**, 151–184 (1983).